

# QUADRATIC EIGENVALUE PROBLEMS: SOLVENTS AND INVERSE PROBLEMS

Peter Lancaster, University of Calgary  
and  
Francoise Tisseur, University of Manchester

Spring, 2009.

## Early Gohberg/Lancaster/Rodman Theory

U. of Calgary, Dept of Math and Stat, Research report 419, 1979. (126pp)

Papers in LAA, Canadian J.Math., IEOT, SIMAX, LAMA. 1977-1982.

Gohberg, I., Lancaster, P., and Rodman, L., *Spectral analysis of matrix polynomials*, Annals of Math, 1982.

Gohberg, I., Lancaster, P., and Rodman, L., *Matrix Polynomials*, Academic Press, 1982. (SIAM 2009 - to appear)

# The quadratic problem

Let  $M, D, K \in \mathbb{C}^{n \times n}$ , all **Hermitian**, with  $M > 0$  and

$$L(\lambda) := M\lambda^2 + D\lambda + K.$$

We will assume that  $L$  is **monic**;  $M = I$ .

We will also consider the important special case of  $D$  and  $K$  **real and symmetric**.

## Factorization of $L(\lambda)$

In general, if  $D^* = D$  and  $K^* = K$ , do there exist matrices  $A$  (a right divisor) and  $S$  (a left divisor) such that

$$I_n \lambda^2 + D \lambda + K = (I_n \lambda - S)(I_n \lambda - A)?$$

## Factorization of $L(\lambda)$

In general, if  $D^* = D$  and  $K^* = K$ , do there exist matrices  $A$  (a right divisor) and  $S$  (a left divisor) such that

$$I_n \lambda^2 + D\lambda + K = (I_n \lambda - S)(I_n \lambda - A)?$$

**YES!** But, existence theorem “evenly divides” eigenvalues with respect to the real axis.

There are many more options for the right and left divisors.

We take advantage of factorizations to solve an inverse problem as follows:

- Assign  $n$  eigenvalues and eigenvectors as those of a right divisor  $A$ .
- Determine a class of compatible left divisors  $S$ .
- Choose a suitable  $S$

## Preliminaries

Suppose that there are  $2r$  **real** evs and  $2(n - r)$  **non-real** ev (in  $n - r$  conjugate pairs).

A right divisor immediately determines  $n$  of these ev's and  $n$  associated **right** eigenvectors, because

$$(I\lambda - A)x = 0, \quad \Rightarrow \quad L(\lambda)x = (I_n\lambda - S)(I_n\lambda - A)x = 0.$$

## Preliminaries

Suppose that there are  $2r$  **real** evs and  $2(n - r)$  **non-real** ev (in  $n - r$  conjugate pairs).

A right divisor immediately determines  $n$  of these ev's and  $n$  associated **right** eigenvectors, because

$$(I\lambda - A)x = 0, \quad \Rightarrow \quad L(\lambda)x = (I_n\lambda - S)(I_n\lambda - A)x = 0.$$

**THE TRIVIAL SOLUTION:** We can solve the **Hermitian** problem by taking  $S = A^*$ . **We eschew this solution!** Why?

Because **real** eigenvalues of  $(I_n\lambda - A^*)(I_n\lambda - A)$  are necessarily **defective** and

$$(I_n\lambda - A^*)(I_n\lambda - A) \geq 0 \quad \text{for all } \lambda \in \mathbb{R}.$$

## The Hermitian case

Let  $S = S_R + iS_I$  where  $S_R = \frac{1}{2}(S + S^*)$  and  $S_I = \frac{-i}{2}(S - S^*)$  are Hermitian. It is not difficult to see that, for a left solvent  $S$ ,

$$S_R A - A^* S_R = \frac{1}{2}(A^2 - (A^*)^2). \quad (1)$$



## The Hermitian case

Let  $S = S_R + iS_I$  where  $S_R = \frac{1}{2}(S + S^*)$  and  $S_I = \frac{-i}{2}(S - S^*)$  are Hermitian. It is not difficult to see that, for a left solvent  $S$ ,

$$S_R A - A^* S_R = \frac{1}{2}(A^2 - (A^*)^2). \quad (1)$$

### Theorem

*Given a matrix  $A \in \mathbb{C}^{n \times n}$  a matrix  $S \in \mathbb{C}^{n \times n}$  is such that both  $S + A$  and  $SA$  are Hermitian if and only if*

$$S = S_R - \frac{1}{2}(A - A^*) = S_R - iA_I,$$

*where  $S_R$  is an Hermitian solution of (1).*

When the proposition holds we have, of course,

$$D = -(S + A) = -(S_R + A_R), \quad K = SA.$$

## The real-symmetric case

With  $A \in \mathbb{R}^{n \times n}$  write

$$A = A_1 + A_2 \quad \text{where} \quad A_1 = \frac{1}{2}(A + A^T), \quad A_2 = \frac{1}{2}(A - A^T),$$

and similarly for  $S$ .

### Corollary

*Given a matrix  $A \in \mathbb{R}^{n \times n}$  a matrix  $S \in \mathbb{R}^{n \times n}$  is such that both  $S + A$  and  $SA$  are real and symmetric if and only if  $S_2 = -A_2$  and  $S_1$  is a real symmetric solution of*

$$S_1 A - A^T S_1 = \frac{1}{2}(A^2 - (A^T)^2). \quad (2)$$

## Real systems in real arithmetic

$A$  is now to be a **real** right divisor and we seek **real** left divisors  $S$ . For the spectrum of  $A$  write

$$\sigma(A) = \{\sigma_1, \dots, \sigma_{2r}\} \cup \{\rho_1, \dots, \rho_s\} \quad (3)$$

where  $2r + s = n$  and

- $\sigma_1 = \mu_1 + i\omega_1, \sigma_2 = \mu_2 - i\omega_2, \dots, \sigma_{2r} = \mu_{2r} - i\omega_{2r}$ , are distinct non-real numbers.
- $\rho_1, \dots, \rho_s$  are distinct real numbers.

## Real systems in real arithmetic

$A$  is now to be a **real** right divisor and we seek **real** left divisors  $S$ . For the spectrum of  $A$  write

$$\sigma(A) = \{\sigma_1, \dots, \sigma_{2r}\} \cup \{\rho_1, \dots, \rho_s\} \quad (3)$$

where  $2r + s = n$  and

- $\sigma_1 = \mu_1 + i\omega_1, \sigma_2 = \mu_2 - i\omega_2, \dots, \sigma_{2r} = \mu_{2r} - i\omega_{2r}$ , are distinct non-real numbers.
- $\rho_1, \dots, \rho_s$  are distinct real numbers.

Consider a **real** Jordan canonical form,  $J_R$ , for  $A$ , assuming that all eigenvalues are simple:

$$J_R = \text{diag} \left[ \left[ \begin{array}{cc} \mu_1 & \omega_1 \\ -\omega_1 & \mu_1 \end{array} \right], \dots, \left[ \begin{array}{cc} \mu_r & \omega_r \\ -\omega_r & \mu_r \end{array} \right], \rho_1, \dots, \rho_s \right], \quad (4)$$

and there is real (e-vector) matrix  $B$  such that  $A = BJ_R B^{-1}$ .

## Real systems in real arithmetic

### Lemma

If  $A$  has  $n$  distinct eigenvalues (as in (3)) and  $B^{-1}AB = J_R$ , then  $S_1$  is a real symmetric solution of  $ZA - A^T Z = 0$  if and only if  $S_1 = B^{-T}FB^{-1}$  where

$$F = \text{diag} \left[ \begin{bmatrix} a_1 & b_1 \\ b_1 & -a_1 \end{bmatrix}, \dots, \begin{bmatrix} a_r & b_r \\ b_r & -a_r \end{bmatrix}, c_1, \dots, c_s \right], \quad (5)$$

and the parameters  $a_j$ ,  $b_j$ ,  $c_k$  are real.

Notice that  $F^T = F$  and  $FJ_R = J_R^T F = (FJ_R)^T$ .

# Real systems in real arithmetic

## Theorem

Given a matrix  $A \in \mathbb{R}^{n \times n}$  and a real matrix  $B$  for which  $A = BJ_R B^{-1}$  and (4) holds, all real matrices  $S$  for which  $A + S$  and  $AS$  are real and symmetric have the form  $S = S_1 + S_2$  where

$$S_1 = A_1 + B^{-T} F B^{-1}, \quad S_2 = -A_2 \quad (6)$$

and  $F$  is an arbitrary real symmetric matrix with the structure of (5).

## Real systems in real arithmetic

Real symmetric system coefficients generated in this way are:

$$D = -(S + A) = -2A_1 - B^{-T}FB^{-1}, \quad (7)$$

$$K = SA = A^T A + B^{-T}(FJ_R)B^{-1}. \quad (8)$$

Clearly, they are determined by the  $n$  real parameters defining  $F$  in (5).

In this construction the number of real (or non-real) eigenvalues in  $A$  can be doubled (in  $L(\lambda)$ ), and there is no further constraint on their positions in the complex plane.

## Real systems in real arithmetic

**Example:** Assign

$$A = BJ_R B^{-1} = \begin{bmatrix} 2 & -2 & 1 & -4 \\ 2 & -3 & 1 & -3 \\ 2 & -2 & -1 & -2 \\ 4 & -3 & 1 & -5 \end{bmatrix}$$

with eigenvalues  $-1 \pm i$ ,  $-2 - 3$ .

If we take

$$F = \text{diag} \left[ \begin{bmatrix} -2 & 0 \\ 0 & 2 \end{bmatrix}, -4, -4 \right],$$

we obtain a matching left divisor

$$S = \begin{bmatrix} -4 & 4 & 1 & -4 \\ 1 & 0 & 1 & -5 \\ 0 & 0 & -9 & 10 \\ 0 & -2 & 13 & -22 \end{bmatrix}$$

with ev  $-28.9022$ ,  $1.5426$ ,  $-4.8432$ ,  $-2.7971$ .



The real symmetric system coefficients obtained are

$$D = \begin{bmatrix} 6 & -3 & -1 & 4 \\ & 0 & -1 & 5 \\ & & 10 & -11 \\ & & & 26 \end{bmatrix}, \quad K = \begin{bmatrix} 7 & -8 & -3 & 13 \\ & 8 & 2 & -8 \\ & & 24 & -41 \\ & & & 99 \end{bmatrix}.$$

## The Hermitian case

As we are interested in the case of mixed, real and non-real eigenvalues, suppose that

$$\sigma(A) = \{\sigma_1, \dots, \sigma_r\} \cup \{\rho_1, \dots, \rho_s\} \quad (9)$$

where  $r + s = n$  and

- $\sigma_1, \dots, \sigma_r$  are distinct non-real numbers,
- $\sigma_j \neq \bar{\sigma}_k$  for  $j, k = 1, \dots, r$  (no conjugate pairs),
- $\rho_1, \dots, \rho_s$  are distinct real numbers.

$$\Delta := \text{diag}(\sigma_1 \cdots \sigma_r), \quad R := \text{diag}(\rho_1 \cdots \rho_s). \quad (10)$$

## The Hermitian case

Then  $A$  has the spectral decomposition

$$A = \sum_{j=1}^r \sigma_j P_j + \sum_{k=1}^s \rho_k Q_k, \quad (11)$$

where  $P_1, \dots, P_r, Q_1, \dots, Q_s$  form a mutually biorthogonal system of rank-one projectors onto the eigenspaces of  $A$  corresponding to the  $\sigma_j, \rho_k$ , respectively.

(If  $r > 0$  then these assumptions do not admit the choice of real matrices  $A$  with non-real eigenvalues.)

# The Hermitian case

## Theorem

Given a matrix  $A \in \mathbb{C}^{n \times n}$  for which (11) and (10) hold, all matrices  $S$  for which  $A + S$  and  $AS$  are Hermitian have the form  $S = S_R + iS_I$  where, for some diagonal  $E \in \mathbb{R}^{s \times s}$ ,

$$S_R = \frac{1}{2}(A + A^*) + T^{-*} \begin{bmatrix} 0 & 0 \\ 0 & E \end{bmatrix} T^{-1}, \quad S_I = \frac{i}{2}(A - A^*). \quad (12)$$

This shows that  $S$  is determined by  $A$  and the  $s$  real parameters defining  $E$ .

(If  $A$  has **no** real eigenvalues, then the last term in the equation for  $S_R$  does not appear and the trivial solution

$$S = \frac{1}{2}(A + A^*) + i \cdot \frac{i}{2}(A - A^*) = A^*$$

is unique.)

## Sign characteristics

In both cases (real/symm. and Hermitian) we can keep track of the sign characteristics of real eigenvalues.

## Hermitian systems with $A \in \mathbb{R}^{n \times n}$

Let the right divisor  $l\lambda - A$  be defined by

$$A = \begin{bmatrix} 1 & 1 & 0 \\ 2 & 3 & 2 \\ 1 & 1 & 2 \end{bmatrix}.$$

Ev of  $A$  are 1, 2, 3.

**(a)** If we choose  $e_1 = e_2 = e_3 = 1$ , then

$$S = \begin{bmatrix} 1 & 2 & 1 \\ -1 & 3 & 1 \\ 0 & 2 & 2 \end{bmatrix} + \sum_{j=1}^3 e_j Y_j = \begin{bmatrix} 3 & 5/2 & 3 \\ -1/2 & 7/2 & 1 \\ 2 & 2 & 5 \end{bmatrix}.$$

$L(\lambda)$  has ev 1, 2, 3 with sign characteristic -1.

The ev of  $S$  (1.78..., 2.69..., 7.02..) interlace those of  $A$  and have sign characteristic +1.

(b) If  $e_1 = e_2 = e_3 = 10$ , then

$$S = \begin{bmatrix} 21 & 7 & 11 \\ 4 & 8 & 1 \\ 20 & 2 & 32 \end{bmatrix}$$

with ev 42.83..., 12.91..., 5.25..., all of +ve type.

This determines a *hyperbolic* system since all ev are real and, assuming the ev's are ordered,  $L(\lambda)$  has  $n$  consecutive ev's of one type followed by  $n$  consecutive ev's of the other type with a gap between the  $n$ th and  $n + 1$ st ev's.

(c) If we let  $e_1 = e_2 = 1, e_3 = -1$ , then

$$S = \begin{bmatrix} 1 & 3/2 & 1 \\ -3/2 & 3 & 0 \\ 0 & 1 & 3 \end{bmatrix}$$

with evs 1.82..., and a conjugate pair:  $2.58.. \pm i(1.05..)$ .

## EPILOGUE: Self-adjoint Jordan triples

$n \times n$  Hermitian systems:  $L(\lambda) = \sum_{j=0}^l A_j \lambda^j$ ,  $A_l > 0$ .

Spectral properties encapsulated in a **self-adjoint Jordan triple**:

$$(X, J, PX^*).$$

$X$  is  $n \times ln$  **complex** (defined by eigenvectors):

$J$  is  $ln \times ln$  **complex** Jordan:

$P$  is  $ln \times ln$  real: all entries 0 or  $\pm 1$ .

(Accounts for sign characteristics of real eigenvalues.)



## EPILOGUE: Self-adjoint Jordan triples

$n \times n$  Hermitian systems:  $L(\lambda) = \sum_{j=0}^l A_j \lambda^j$ ,  $A_l > 0$ .

Spectral properties encapsulated in a **self-adjoint Jordan triple**:

$$(X, J, PX^*).$$

$X$  is  $n \times ln$  **complex** (defined by eigenvectors):

$J$  is  $ln \times ln$  **complex** Jordan:

$P$  is  $ln \times ln$  real: all entries 0 or  $\pm 1$ .

(Accounts for sign characteristics of real eigenvalues.)

$\Rightarrow$  **Hermitian** moments  $\Gamma_j = X(J^j)PX^*$ ,  $j = 0, 1, \dots$

$\Rightarrow$  formulae for **Hermitian** coefficients  $A_j$ .

# Real symmetric systems

Are Hermitian! But have **more** structure.

Spectral properties encapsulated in a **real self-adjoint Jordan triple**:

$$(X_R, J_R, P_R X_R^T).$$

$X_R$  is  $n \times ln$  **real** (defined by real and im. parts of eigenvectors):

$J_R$  is  $ln \times ln$  **real** Jordan:

$P_R$  is  $ln \times ln$  real: all entries 0 or  $\pm 1$ .

(Essentially unchanged.)

# Real symmetric systems

Are Hermitian! But have **more** structure.

Spectral properties encapsulated in a **real self-adjoint Jordan triple**:

$$(X_R, J_R, P_R X_R^T).$$

$X_R$  is  $n \times ln$  **real** (defined by real and im. parts of eigenvectors):

$J_R$  is  $ln \times ln$  **real** Jordan:

$P_R$  is  $ln \times ln$  real: all entries 0 or  $\pm 1$ .

(Essentially unchanged.)

$\Rightarrow$  **real symmetric** moments  $\Gamma_j = X(J^j)PX^*$ ,  $j = 0, 1, \dots$

$\Rightarrow$  formulae for **real symmetric** coefficients  $A_j$ .



Time to go!