

# Hermitian Matrix Polynomials with Real Eigenvalues

**Françoise Tisseur**  
**School of Mathematics**  
**The University of Manchester**

`ftisseur@ma.man.ac.uk`  
`http://www.ma.man.ac.uk/~ftisseur/`

Joint work with **Maha Al-Ammari**  
**MOPNET II, March 29-30, 2010**

# Hermitian Eigenvalue Problem

Consider

$$Ax = \lambda x, \quad A = A^* \in \mathbb{C}^{n \times n}$$

- ▶ Many desirable properties. In particular,
  - real eigenvalues,
  - diagonalizable by congruences,
  - well-conditioned eigenvalues.
  
- ▶ Variety of special algorithms.

# Hermitian Eigenvalue Problem

Consider

$$Ax = \lambda x, \quad A = A^* \in \mathbb{C}^{n \times n}$$

- ▶ Many desirable properties. In particular,
  - real eigenvalues,
  - diagonalizable by congruences,
  - well-conditioned eigenvalues.
  
- ▶ Variety of special algorithms.

**What are the closest analogues of this class of problems for Hermitian matrix polynomials?**

# Hermitian Eigenproblem

We consider

- ▶ Generalized eigenvalue problem:  $L(\lambda)x = 0$ ,

$$L(\lambda) = \lambda A - B, \quad A = A^*, \quad B = B^*.$$

- ▶ Polynomial eigenvalue Problem:  $P(\lambda)x = 0$ ,

$$P(\lambda) = \sum_{j=0}^m \lambda^j A_j, \quad A_j = A_j^*, \quad j = 0: m.$$

- $mn$  e'vals, all finite when  $A_m$  nonsingular,
- $\infty$  and  $0$  e'vals when  $A_m$  and  $A_0$  singular, resp.,
- $\Lambda(P)$  is symmetric with respect to the real axis.

# Hermitian Pencils/Polynomials

The following are known to have **real e'vals**:

- definite pencils,
- definitizable pencils,
- overdamped quadratics,
- gyroscopically stabilized quadratics
- hyperbolic matrix polynomials,
- quasihyperbolic matrix polynomials,
- definite matrix polynomials.

Another common feature: their **e'vals are all of definite type**.

# Eigenvalue Types

A finite real e'val  $\lambda_0$  of  $P(\lambda)$  Hermitian is of


- **positive type** if  $x^* P'(\lambda_0)x > 0$  for all  $0 \neq x \in \ker P(\lambda_0)$ ,
- **negative type** if  $x^* P'(\lambda_0)x < 0$  for all  $0 \neq x \in \ker P(\lambda_0)$ .
- **definite type** if it is either of positive or negative type.

# Eigenvalue Types

A finite real e'val  $\lambda_0$  of  $P(\lambda)$  Hermitian is of


- **positive type** if  $x^* P'(\lambda_0)x > 0$  for all  $0 \neq x \in \ker P(\lambda_0)$ ,
- **negative type** if  $x^* P'(\lambda_0)x < 0$  for all  $0 \neq x \in \ker P(\lambda_0)$ .
- **definite type** if it is either of positive or negative type.

$$q_x(\lambda) = x^* P(\lambda)x$$



negative type

$$q_x(\lambda) = x^* P(\lambda)x$$



positive type

# Examples

- Simple e'vals are always of definite type.
- The pencil  $L(\lambda) = \lambda \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} + \begin{bmatrix} -\alpha & 0 \\ 0 & \alpha \end{bmatrix}$  has a semisimple e'val  $\lambda_0 = \alpha$  with e'vecs  $\mathbf{e}_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ ,  $\mathbf{e}_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$ , and

$$\mathbf{e}_1^* L'(\alpha) \mathbf{e}_1 = 1, \quad \mathbf{e}_2^* L'(\alpha) \mathbf{e}_2 = -1.$$

$\Rightarrow \lambda_0 = \alpha$  is of **mixed type**.



# Eigenvalue Type at $\infty$

$$\text{rev}P(\lambda) := \lambda^m P(1/\lambda) = \lambda^m A_0 + \lambda^{m-1} A_1 + \cdots + \lambda A_{m-1} + A_m.$$

$\lambda = \infty$  is an e'val of  $P(\lambda)$  iff 0 is an e'val of  $\text{rev}P(\lambda)$ .

Can show that

$$x^* P'(\lambda_0) x = -\lambda_0^{m-2} x^* (\text{rev}P)'(1/\lambda_0) x, \quad \lambda_0 \neq 0.$$

# Eigenvalue Type at $\infty$

$$\text{rev}P(\lambda) := \lambda^m P(1/\lambda) = \lambda^m A_0 + \lambda^{m-1} A_1 + \cdots + \lambda A_{m-1} + A_m.$$

$\lambda = \infty$  is an e'val of  $P(\lambda)$  iff 0 is an e'val of  $\text{rev}P(\lambda)$ .

Can show that

$$x^* P'(\lambda_0) x = -\lambda_0^{m-2} x^* (\text{rev}P)'(1/\lambda_0) x, \quad \lambda_0 \neq 0.$$

$\lambda_0 = \infty$  as an e'val of  $P$  is of

- positive type if  $x^* A_{m-1} x < 0$  for all  $0 \neq x \in \ker A_m$ ,
- negative type if  $x^* A_{m-1} x > 0$  for all  $0 \neq x \in \ker A_m$ .

# Systems of Differential Equations

The solutions to

$$\sum_{j=0}^m i^j A_j \frac{d^j u(t)}{dt^j} = 0, \quad t \in \mathbb{R},$$

are bounded on  $[0, \infty)$  iff  $P(\lambda) = \sum_{j=0}^m \lambda^j A_j$ ,  $\det(A_m) \neq 0$  has real semisimple e'vals.

Solutions remain bounded under small perturbations of  $A_j$  iff the e'vals of  $P$  are real and of definite type [Gohberg, Lancaster, Rodman 82].

# Quasidefinite Matrix Polynomials

## Definition (Al-Ammari, T., 10)

A Hermitian matrix polynomial  $P$  is **quasidefinite** if

- $\Lambda(P) \subset \mathbb{R} \cup \{\infty\}$ , and
- all e'vals are of definite type.

# Quasidefinite Matrix Polynomials

## Definition (Al-Ammari, T., 10)


A Hermitian matrix polynomial  $P$  is **quasidefinite** if

- $\Lambda(P) \subset \mathbb{R} \cup \{\infty\}$ , and
- all e'vals are of definite type.

Quasidefinite polynomials include

- definite pencils,
- definitizable pencils,
- overdamped quadratics,
- gyroscopically stabilized quadratics,
- hyperbolic matrix polynomials,
- quasihyperbolic matrix polynomials,
- definite matrix polynomials.

# Classification of Quasidefinite Polynomials




# Definite Pencils $L(\lambda) = \lambda A - B$

$L(\lambda)$  is **definite** if it satisfies any one of (P1), (P2), (P3).

## Theorem

*The following are equivalent:*


- (P1)  $\Lambda(L) \subset \mathbb{R} \cup \{\infty\}$  with all e'vals of definite type and e'vals of +ve type are separated from the e'vals of -ve type.
- (P2) The matrix  $L(\mu)$  is definite for some  $\mu \in \mathbb{R} \cup \{\infty\}$ .
- (P3) For every nonzero vector  $x$ , the scalar equation  $x^* L(\lambda)x = 0$  has exactly one zero in  $\mathbb{R} \cup \{\infty\}$ .



# Homogeneous Form

$$P(\alpha, \beta) = \sum_{j=0}^m \alpha^j \beta^{m-j} A_j, \quad L(\alpha, \beta) = \alpha A - \beta B.$$

- ▶ E'val  $\lambda$  identified with any pair  $(\alpha, \beta) \neq (0, 0)$  s.t.  $\lambda = \alpha/\beta$ .
- ▶  $\lambda = 0$  represented by  $(0, \beta)$ ,  $\lambda = \infty$  represented by  $(\alpha, 0)$ .
- ▶ With  $\alpha^2 + \beta^2 = 1$ , have direct correspondence between  $\lambda \in \mathbb{R} \cup \{\infty\}$  and  $(\alpha, \beta)$  on unit circle:





# Homogeneous Rotation

$\tilde{P}(\tilde{\alpha}, \tilde{\beta})$  is obtained from  $P(\alpha, \beta)$  by **homogenous rotation** if

$$G \begin{bmatrix} \alpha \\ \beta \end{bmatrix} = \begin{bmatrix} c & s \\ -s & c \end{bmatrix} \begin{bmatrix} \alpha \\ \beta \end{bmatrix} =: \begin{bmatrix} \tilde{\alpha} \\ \tilde{\beta} \end{bmatrix}, \quad c, s \in \mathbb{R}, \quad c^2 + s^2 = 1$$

and

$$P(\alpha, \beta) = \sum_{j=0}^m \alpha^j \beta^{m-j} A_j = \sum_{j=0}^m \tilde{\alpha}^j \tilde{\beta}^{m-j} \tilde{A}_j := \tilde{P}(\tilde{\alpha}, \tilde{\beta}).$$


- ▶ E'vecs remain the same but e'vals are rotated.
- ▶  $\tilde{A}_m = P(c, s)$ .
- ▶ Use rotation  $G$  to transform  $P$  with  $\det(A_m) = 0$  or  $A_m$  indefinite to  $\tilde{P}$  with  $\tilde{A}_m$  nonsingular or  $\tilde{A}_m > 0$ .

# Homogeneous Rotation and Types

Let  $\lambda_j$  be an e'val of  $P$  rotated to  $\tilde{\lambda}_j$ . E'val types are related by:

$$x^* \tilde{P}'_{\tilde{\lambda}}(\tilde{\lambda}_j) x = (c - \lambda_j s)^{m-2} \cdot x^* P'_{\lambda}(\lambda_j) x \quad \text{if } \lambda_j, \tilde{\lambda}_j \text{ are finite.}$$

$c - \lambda_j s = \det \begin{bmatrix} c & \lambda_j \\ s & 1 \end{bmatrix} > 0$  if  $\lambda_j = (\lambda_j, 1)$  that lies counterclockwise from  $(c, s)$ .



# Hyperbolic Polynomials $P(\lambda) = \sum_{j=0}^m \lambda^j A_j$

$P(\lambda)$   $n \times n$  and Hermitian is **hyperbolic** if it satisfies any one of (P1) [Al-Ammari, T. 10], (P2), (P3) [Markus 88]. .

## Theorem

*The following are equivalent:*

(P1) *All e'vals are real, finite, of definite type, and s.t.*

$$\underbrace{\lambda_{mn} \leq \dots \leq \lambda_{(m-1)n+1}}_{(-1)^{m-1} \text{ type}} < \dots < \underbrace{\lambda_{2n} \leq \dots \leq \lambda_{n+1}}_{\text{negative type}} < \underbrace{\lambda_n \leq \dots \leq \lambda_1}_{\text{positive type}}.$$

(P2) *There exist  $\mu_j \in \mathbb{R} \cup \{\infty\}$  s.t.  $\infty = \mu_0 > \mu_1 > \dots > \mu_{m-1}$ ,*

$$(-1)^j P(\mu_j) > 0, \quad j = 0: m-1.$$

(P3)  *$A_m > 0$  and for every nonzero  $x \in \mathbb{C}^n$ , the scalar equation  $x^* P(\lambda)x = 0$  has  $m$  distinct real and finite zeros.*

# Acoustic Fluid-structure Interaction Problem

Consider generalized eigenproblem

$$\omega \begin{bmatrix} M_s & 0 \\ M_{fs} & M_f \end{bmatrix} + \begin{bmatrix} K_s & -M_{fs}^* \\ 0 & K_f \end{bmatrix},$$

where  $0 < M_s, K_s \in \mathbb{C}^{n \times n}$  and  $0 < M_f, K_f \in \mathbb{C}^{m \times m}$ .

Multiplying 1st block row by  $-\omega$  yields Hermitian quadratic

$$Q(\omega) = \omega^2 \begin{bmatrix} -M_s & 0 \\ 0 & 0 \end{bmatrix} + \omega \begin{bmatrix} -K_s & M_{fs}^* \\ M_{fs} & M_f \end{bmatrix} + \begin{bmatrix} 0 & 0 \\ 0 & K_f \end{bmatrix}.$$

- ▶  $Q$  is not hyperbolic,
- ▶  $\text{rev}Q(\omega) = \omega^2 Q(1/\omega)$  is not hyperbolic.
- ▶ However,  $Q$  is a definite polynomial .

# Definitizable Pencils

Definition: An  $n \times n$  Hermitian pencil  $L(\lambda) = \lambda A - B$  is **definitizable** if it satisfies any one of (P1), (P2), (P3).

## Theorem

*The following are equivalent:*

- (P1) *L has real, finite e'vals of definite type.*
- (P2)  *$\det(A) \neq 0$  and there exists a real polynomial  $q$  s.t.  $Aq(A^{-1}B) > 0$ .*
- (P3)  *$\det(A) \neq 0$  and the scalar equation  $x^* L(\lambda)x = 0$  has one zero in  $\mathbb{R}$  for all e'vecs  $x \in \mathbb{C}^n$  of  $L$ .*

Proofs in [Lancaster, Ye, 93], except (P3).

# Saddle Point Problems

Want to solve large linear systems  $\mathcal{A}x = b$  with

$$\mathcal{A} = \begin{bmatrix} A & B \\ B & -C \end{bmatrix},$$

where  $A = A^T \in \mathbb{R}^{n \times n}$ ,  $A > 0$  and  $C = C^T \in \mathbb{R}^{m \times m}$ ,  $C \geq 0$ .

$\mathcal{A}$  is **indefinite**: it has  $n$  positive e'vals and  $\text{rank}(C + BA^{-1}B^T)$  negative e'vals.

# Saddle Point Problems

Want to solve large linear systems  $\mathcal{A}x = b$  with

$$\mathcal{A} = \begin{bmatrix} A & B \\ B & -C \end{bmatrix},$$

where  $A = A^T \in \mathbb{R}^{n \times n}$ ,  $A > 0$  and  $C = C^T \in \mathbb{R}^{m \times m}$ ,  $C \geq 0$ .

$\mathcal{A}$  is **indefinite**: it has  $n$  positive e'vals and  $\text{rank}(C + BA^{-1}B^T)$  negative e'vals.

If  $\lambda \mathcal{J} - \mathcal{A}$  is definitizable with  $\mathcal{J} = \begin{bmatrix} I_n & 0 \\ 0 & -I_m \end{bmatrix}$  then there exists a well-define CG method for solving linear systems with  $\mathcal{J}\mathcal{A}$  (see [Liesen & Parlett 08]).

( $\mathcal{J}\mathcal{A}$  is  $\mathcal{J}q(\mathcal{J}\mathcal{A})$  symmetric for some  $q$  s.t.  $\mathcal{J}q(\mathcal{J}\mathcal{A}) > 0$ .)

# Hermitian Linearizations

Let  $\mathbb{L}_1(P) = \{ L(\lambda) : L(\lambda)(\Lambda \otimes I_n) = v \otimes P(\lambda), v \in \mathbb{C}^m \}$ ,

where  $\Lambda = [\lambda^{m-1}, \lambda^{m-2}, \dots, 1]^T \in \mathbb{C}^m$ .

$$\begin{aligned} \mathbb{H}(P) &:= \{ L(\lambda) = \lambda A - B \in \mathbb{L}_1(P) : A^* = A, B^* = B \} \\ &= \left\{ \sum_{j=1}^m v_j (\lambda B_j - B_{j-1}), v \in \mathbb{R}^m \right\}, \end{aligned}$$

where  $B_j$  is a direct sum of block Hankel matrices.

Almost all pencils in  $\mathbb{H}(P)$  are Hermitian linearizations of  $P$ .

**Do they preserve additional properties?**



# Linearizations and E'val Types

For an eigenpair  $(\lambda_0, x)$  of  $P$  and  $L(\lambda) \in \mathbb{H}(P)$  with vector  $v$ , we have

$$z^* L'(\lambda_0) z = \Lambda_0^T v \cdot x^* P'(\lambda_0) x,$$

where  $(z, \lambda_0)$  is an eigenpair of  $L$ ,  $\Lambda_0 = [\lambda_0^{m-1}, \lambda_0^{m-2}, \dots, 1]^T$ .

# Linearizations and E'val Types

For an eigenpair  $(\lambda_0, x)$  of  $P$  and  $L(\lambda) \in \mathbb{H}(P)$  with vector  $v$ , we have

$$z^* L'(\lambda_0) z = \Lambda_0^T v \cdot x^* P'(\lambda_0) x,$$

where  $(z, \lambda_0)$  is an eigenpair of  $L$ ,  $\Lambda_0 = [\lambda_0^{m-1}, \lambda_0^{m-2}, \dots, 1]^T$ .

## Theorem

- $P$  is quasihyperbolic iff any  $L \in \mathbb{H}(P)$  is definitizable [Al-Ammari, T., 10].
- $P$  is definite iff  $P$  has a definite linearization  $L \in \mathbb{H}(P)$ . [Higham, Mackey, T. 09].
- $P$  is hyperbolic iff  $P$  has a definite linearization  $\lambda A - B \in \mathbb{H}(P)$  with  $A$  definite.

# Diagonalizable Pencils

Hermitian pencils are diagonalizable by congruence iff e'vals belong to  $\mathbb{R} \cup \{\infty\}$  and are semisimple (see [Lancaster, Rodman 05]).

- ▶ Definite pencils are diagonalizable.
- ▶ Definitizable pencils are diagonalizable.

# Diagonalizable Pencils

Hermitian pencils are diagonalizable by congruence iff e'vals belong to  $\mathbb{R} \cup \{\infty\}$  and are semisimple (see [Lancaster, Rodman 05]).

- ▶ Definite pencils are diagonalizable.
- ▶ Definitizable pencils are diagonalizable.

**What can we say about (quasi)hyperbolic and definite matrix polynomials?**

# Strictly Isospectral Polynomials

$P$  is **isospectral** to  $\hat{P}$  if  $\Lambda(P) = \Lambda(\hat{P})$  with same partial multiplicities.

$P$  and  $\hat{P}$  are **strictly isospectral** if they are isospectral and share the **same sign characteristic**.

# Strictly Isospectral Polynomials

$P$  is **isospectral** to  $\widehat{P}$  if  $\Lambda(P) = \Lambda(\widehat{P})$  with same partial multiplicities.

$P$  and  $\widehat{P}$  are **strictly isospectral** if they are isospectral and share the **same sign characteristic**.

Let  $P, \widehat{P}$  be quasihyperbolic and strictly isospectral and let  $L \in \mathbb{H}(P), \widehat{L} \in \mathbb{H}(\widehat{P})$  with vector  $v$ .

There exist nonsingular  $X, \widehat{X}$  s.t.

$$XL(\lambda)X^* = \lambda \begin{bmatrix} I_k & 0 \\ 0 & -I_{n-k} \end{bmatrix} - \begin{bmatrix} J_+ & 0 \\ 0 & -J_- \end{bmatrix} = \widehat{X}\widehat{L}(\lambda)\widehat{X}^*.$$

$\widehat{X}^{-1}X$  defines a **structure preserving congruence**.

# Diagonalizable by SPC

Definition:  $P(\lambda)$ , Hermitian and of degree  $m$  is **diagonalizable by structure preserving congruence** (SPC) if it is strictly isospectral to a real diagonal matrix polynomial of degree  $m$ .

- ▶ Quasidefinite quadratics are always strictly isospectral to quasidefinite diagonal quadratics.
- ▶ Definite matrix polynomial are always strictly isospectral to definite diagonal matrix polynomials.

# Diagonalizable by SPC

Definition:  $P(\lambda)$ , Hermitian and of degree  $m$  is **diagonalizable by structure preserving congruence** (SPC) if it is strictly isospectral to a real diagonal matrix polynomial of degree  $m$ .

## Theorem (Al-Ammari, T. 10)

*An  $n \times n$  quasihyperbolic matrix polynomial of degree  $m$  is diagonalizable by SPC iff there is a grouping of its  $e$ 'vals and their types into  $n$  subsets of  $m$  distinct  $e$ 'vals, which when ordered have alternating types.*

- ▶ Quasidefinite quadratics are always strictly isospectral to quasidefinite diagonal quadratics.
- ▶ Definite matrix polynomial are always strictly isospectral to definite diagonal matrix polynomials.



# Concluding Remarks

- ▶ Gave a unified treatment of the many subclasses of Hermitian matrix polynomials with real eigenvalues.
- ▶ Identified classes of Hermitian matrix polynomials that are diagonalizable by SPC.
- ▶ Results useful in the solution of the inverse problem.
- ▶ Investigate analogous results for palindromic and odd/even matrix polynomials.

For paper see:

M. Al-Ammari and F. Tisseur. *Hermitian Matrix Polynomials with Real Eigenvalues of Definite Type. Part I: Classification*, MIMS EPrint 2010.9, The University of Manchester, 2010.